

Heilpflanzendiversität an Saum- und Ruderalstandorten

Literatur (Gesamtliste)

- AKADEMIEN DER WISSENSCHAFTEN SCHWEIZ (2014): Bienen und andere Bestäuber: Bedeutung für Landwirtschaft und Biodiversität. Swiss Academics Factsheets 9 (1).
- AKGUL, C. & G. SAGLIKOGU (2005): Antibacterial activity of crude methanolic extract and its fractions of aerial parts of *Anthemis tinctoria*. Indian J. Biochem. Biophys. 42: 395–397.
- AL-SNAFI, A.E. (2017): A review on *Erodium cicutarium*: A potential medicinal plant. Indo Am. J. P. Sci. 4: 110–116.
- AN, H. J., JEONG, H. J., UM, J. Y., KIM, H. M. & S. H. HONG (2006): *Glechoma hederacea* inhibits inflammatory mediator release in IFN-gamma and LPS-stimulated mouse peritoneal macrophages. J. Ethnopharmacol. 106: 418–424.
- BAUMBERGER, J. (2010): Phytochemical and biological investigations on *Carduus crispus* L. (Asteraceae). Dipl.-Arb., Graz, 81 S.
- BENARBA, B., MEDDAH, B. & A. AOUES (2012): *Bryonia dioica aqueous* extract induces apoptosis through mitochondrial intrinsic pathway in BL41 Burkitt's lymphoma cells. J. Ethnopharmacol. 141: 510–516.
- BERAN, H. & W. VÖLKL (2011): Der Niedergang der Saumstrukturen. Vogelschutz-Magazin für Arten- und Biotopschutz 3: 14–17.
- BERRINGTON, D. & N. LALL (2012): Anticancer activity of certain herbs and spices on the cervical epithelial carcinoma (HeLa) cell line. Evid. Based Complement. Altern. Med.:doi:10.1155/2012/564927.
- BfN [BUNDESAMT FÜR NATURSCHUTZ] (2003): Floraweb – Daten und Informationen zu Wildpflanzen und zur Vegetation Deutschlands. <http://floraweb.de>, abgerufen am 08.01.2017.
- BIESMEIJER, J.C., ROBERTS, S.P.M., REEMER, M., OHLEMÜLLER, R., EDWARDS, M., PEETERS, T., SCHAFFERS, A.P., POTTS, S.G., KLEUKERS, R. & C.D. THOMAS (2006): Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. Science 313(5785): 351–354.
- BRITTAİN, C., WILLIAMS, N., KREMEN, C. & A.-M. KLEIN (2013): Synergetic effects of non-Apis bees and honey bees for pollination services. Proc. R. Soc. Lond. B Biol. Sci. 280: 1471–2954.
- CHEN, J.J., TSAI, Y.C., HWANG, T.L. & T.C. WANG (2011): Thymol, benzofuranoid, and phenylpropanoid derivatives: anti-inflammatory constituents from *Eupatorium cannabinum*. J. Nat. Prod. 74: 1021–1027.
- CONFORTI, F., PERRI, N., MENICHINI, F., MARELLI, M., UZUNOV, D., STATTI G.A., & F. MENICHINI (2012): Wild mediterranean dietary plants as inhibitors of pancreatic lipase. Phytother. Res. 26: 600–604.
- DACI, A., GOLD-BINDER, M., GARZON, D., PATEA, A. & G. BERETTA (2014): Standardization of solvent extracts from *Onopordum acanthium* fruits by GC-MS, HPLC-UV/DAD, HPLC-TQMS and ¹H-NMR and evaluation of their inhibiting effects on the expression of IL-8 and E-selectin in immortalized endothelial cells (HUVEC tert). Nat. Prod. Commun. 9: 945–948.
- DE LA RÚA, P., JAFFÉ, R., DALL'OLIO, R., MUÑOZ, I. & J. SERRANO (2009): Biodiversity, conservation and current threats to European honeybees. Apidologie 40 (3): 263–284.
- DI PAOLA, R., ESPOSITO, E., MAZZON, E., RICCARDI, L., CAMINITI, R., DAL TOSO, R., PRESSI, G. & S. CUZZOCREA (2009): Teupolioside, a phenylpropanoid glycosides of *Ajuga reptans*, biotechnologically produced by IRBN22 plant cell line, exerts beneficial effects on a rodent model of colitis. Biochem. Pharmacol. 77: 845–857.
- DIB, I., FAUCONNIER, M.-L., SINDIC, M., BELMEKKI, F., ASSAIDI, A., BERRABAH, M., MEKHFI, H., AZIZ, M., LEGSSYER, A., BNOUHAM, M. & A. ZIYYAT (2017): Chemical composition, vasorelaxant, antioxidant and antiplatelet effects of essential oil of *Artemisa campestris* L. from Oriental Morocco. BMC Complement. Altern. Med. 17: 1–15.
- DÜLL, R. & H. KUTZELNIGG (1994): Botanisch-ökologisches Exkursionstaschenbuch. 5. Aufl., Quelle & Meyer, Wiesbaden, 590 S.
- DZHAMBAYOV, B., DASKALOVA, S., MONTEVA, A. & N. POPOV (2002): In vitro screening for antitumor activity of *Clinopodium vulgare* L. (Lamiaceae) extracts. Biol. Pharm. Bull. 25: 499–504.
- FÜLLE, K.-M. (2014): Bienengerechte Fütterung in der Praxis. 21 S.
- GOULSON, D., NICHOLLS, E., BOTÍAS, C. & E. L. ROTHERAY (2015): Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. Science 347. DOI:10.1126/science.1255957.
- HABIBATNI, S., ZOHRA, A.F., KHALIDA, H., ANWAR, S., MANSI, I. & N.A. AWADH ALI (2017): In-vitro antioxidant, xanthine oxidase-inhibitory and in-vivo anti-inflammatory, analgesic, antipyretic activity of *Onopordum acanthium*. Int. J. Phytomed. 9: 92–100.
- HAEUPLER, H. & T. MUER (2000): Bildatlas der Farn- und Blütenpflanzen Deutschlands. Ulmer Verlag, Stuttgart, 759 S.
- HARRISON, P. A., BERRY, P. M., SIMPSON, G., HASLETT, J. R., Blicharska, M., BUCUR, M., DUNFORD, R., EGOH, B., GARCIA-LLORENTE, M., GEAMĂNĂ, GEERTSEMA, W., LOMMELEN, E., MEIRESONNE, L. & F. TURKELBOOM (2014): Linkages between biodiversity attributes and ecosystem services: A systematic review. Ecosystem Services 9: 191–203.
- HOEHN, P., TSCHARNTKE, T., TYLIANAKIS, J. M. & I. STEFFAN-DEWENTER (2008): Functional group diversity of bee pollinators increases crop yield. Proc. R. Soc. B 275: 2283–2291.
- HOLZSCHUH, A., DUDENHÖFER, J.-H. & T. TSCHARNTKE (2012): Landscapes with wild bee habits enhance pollination, fruit set and yield of sweet cherry. Biol. Conserv. 153: 101–107.
- HUYAN, T., LI, Q., WANG, Y.L., LI, J., ZHANG, J.Y., LIU, Y.X., SHAHID, M.R., YANG, H. & H.Q. LI (2016): Anti-tumor effect of hot aqueous extracts from *Sonchus oleraceus* (L.) L. and *Juniperus sabina* L. Two traditional medicinal plants in China. J. Ethnopharmacol. 185: 289–299.
- JÄGER, E.J. (Hrsg.) (2011): Rothmaler. Exkursionsflora von Deutschland. Gefäßpflanzen: Grundband. 20. Aufl. Spektrum Akademischer Verlag, 930 S.
- JAIN, K.A. (2014): Mechanistic approach to determination of anti-diabetic activity of *Calystegia sepium* R.Br. flowering plants in normal and streptozotocin induced rats. Asian J. Res. Pharma. Sci. 4: 55–70.
- JANBAZ, K.H., LATIF, M.F., SAQIB, F., IMRAM, I., ZIA-UL-HAQ, M. & V. DE FEO (2013): Pharmacological effects of *Lactuca serriola* L. in experimental model of gastrointestinal, respiratory, and vascular ailments. Evid. Based Complement. Alternat. Med. 2013: 1–9.
- KENNY, O., SMYTH, T.J., WALSH, D., KELLEHER, C.T., HEWAGE, C.M. & N.P. BRUNTON (2014): Investigating the potential of underutilised plants from the Asteraceae family as a source of natural antimicrobial and antioxidant extracts. Food Chem. 161: 79–86.

- KHAN, H., SAEED, M., GILANI, A. U., KHAN, M. A., KHAN, I. & N. ASHRAF (2011): Antinociceptive activity of aerial parts of *Polygonatum verticillatum*: attenuation of both peripheral and central pain mediators. *Phytother. Res.* 25: 1024–1030.
- KHAN, R. A., KHAN, M. R. & S. SAHREEN (2012): Brain antioxidant markers, cognitive performance and acetylcholinesterase activity of rats: efficiency of *Sonchus asper*. *Behav. Brain Funct.* 8/2012: 1–7.
- KLEIN, A.-M., VAISSIERE, B. E., CANE, J. H., STEFFAN-DEWENTER, I., CUNNINGHAM, S. A., KREMEN, C. & T. TSCHARNTKE (2007): Importance of pollination in changing landscapes for world crops. *PRS-B.* 274(1608): 303–313.
- KLEIJN, D. & I. RAEMAKERS (2008): A retrospective analysis of pollen host plant use by stable and declining bumble bee species. *Ecology* 89: 1811–1823.
- KOKANOVA-NEDIALKOVA, Z., KONDEVA-BURDINA, M., ZHELEVA-DIMITROVA, D., TZANKOVA, V., NIKOLOV, S., HEILMANN, J. & P. T. NEDIALKOV (2015): 6-Methoxyflavonol glycosides with in vitro hepatoprotective activity from *Chenopodium bonus-henricus* roots. *Nat. Prod. Commun.* 10: 1377–1380.
- KOKANOVA-NEDIALKOVA, Z., NEDIALKOV, P. T., KONDEVA-BURDINA, M., SIMEONOVA, R., TZANKOVA, V. & D. ALUANI (2017): *Chenopodium bonus-henricus* L. – A source of hepatoprotective flavonoids. *Fitoterapia* 118: 13–20.
- KOVÁTS, N., ÁCS, A., GÖLÖNCSEI, F. & A. BARABÁS (2011): Quantifying of bactericide properties of medicinal plants. *Plant Signal. Behav.* 6: 777–779.
- LANUV [LANDESAMT FÜR NATUR, UMWELT UND VERBRAUCHERSCHUTZ NORDRHEIN-WESTFALEN] (2017): Blühende Vielfalt am Wegesrand. Praxis-Leitfaden für artenreiche Weg- und Feldraine. LANUV-Info 39.
- LI, J., LI, Q. W., GAO, D. W., HAN, Z. S. & K. LI (2008): Antitumor effects of total alkaloids isolated from *Solanum nigrum* in vitro and in vivo. *Pharmazie* 63: 534–538.
- LIN, H. M., TSENG, H. C., WANG, C. J., LIN, J. J., LO, C. W. & F. P. CHOU (2008): Hepatoprotective effects of *Solanum nigrum* Linn extract against CCl₄-induced oxidative damage in rats. *Chem. Biol. Interact.* 171: 283–293.
- LIN, B., ZHAO, Y., HAN, P., YUE, W., MA, X. Q., RAHMAN, K., ZHENG, C. J., QIN, L. P. & T. HAN (2014): Anti-arthritic activity of *Xanthium strumarium* L. extract on complete Freund's adjuvant induced arthritis in rats. *J. Ethnopharmacol.* 155: 248–255.
- MAHERNIA, S., BAGHERZADEH, K., MOJAB, F. & M. AMANLOU (2015): Urease inhibitory activities of some commonly consumed herbal medicines. *Iran. J. Pharm. Res.*, 14: 943–947.
- MATSUMOTO, T., MORIYA, M., SAKURAI, M. H., KIYOHARA, H., TABUCHI, Y. & H. YAMADA (2008): Stimulatory effect of a peptic polysaccharide from a medicinal herb, the roots of *Bupleurum falcatum* L., on G-CSF secretion from intestinal epithelial cells. *Int. Immunopharmacol.* 8: 581–588.
- MIHAILOVIĆ, V., KREFT, S., BENKOVIĆ, E. T., IVANOVIĆ, N. & M. S. STANKOVIĆ (2016): Chemical profile, antioxidant activity and stability in stimulated gastrointestinal tract model system of three *Verbascum* species. *Ind. Crop. Prod.* 89: 141–152.
- MLINARIC, A., KREFT, S., UMEK, A. & B. STRUKELJ (2000): Screening of selected plant species for in vitro inhibiting activity on HIV-1 reverse transcriptase (HIV-1RT). *Pharmazie* 55: 75–77.
- MOALLEM, S. A., HOSSEINZADEH, H. & F. GHONCHEH (2007): Evaluation of antidepressant effects of aerial parts of *Echium vulgare* on mice. *Iran. J. Basic Med. Sci.* 10: 189–196.
- NAZARUK, J., CZECHOWSKA, S. K., MARKIEWICZ, R. & M. H. BORAWSKA (2008): Polyphenolic compounds and in vitro antimicrobial and antioxidant activity of aqueous extracts from leaves of some *Cirsium* species. *Nat. Prod. Res.* 22: 1583–1588.
- NEITZKE, M. (2005): Einheimische Heilpflanzen (Standort, Wirkung, Geschichte). Vorlesungsskript (unveröff.).
- NEITZKE, M. (2015): Heilpflanzendiversität in den Ackerökosystemen Nordrhein-Westfalens. *Natur in NRW* 4/2015: 32–36.
- NEITZKE, A. (2011): Saumbiotop und Vernetzungsstrukturen in Mais. *Mais* 1: 8–11.
- NIKOLOVA, M. (2011): Screening of radical scavenging activity and polyphenol content of Bulgarian plant species. *Pharmacognosy Res.* 3: 256–259.
- NOLKEMPER, S., REICHLING, J., STINTZING, F. C., CARLE, R. & P. SCHNITZLER (2006): Antiviral effect of aqueous extracts from species of the Lamiaceae family against Herpes simplex virus type 1 and type 2 in vitro. *Planta Med.* 72: 1378–1382.
- OBERDORFER, E. (Hrsg.) (1978): Süddeutsche Pflanzengesellschaften. Sand- und Trockenrasen, Heide und Borstgrasgesellschaften, alpine Magerrasen, Saum-Gesellschaften, Schlag- und Hochstaudenfluren. Teil II. 3. Aufl. Stuttgart, Fischer, 455 S.
- OBERDORFER, E. (1983): Pflanzensoziologische Exkursionsflora. 5. Aufl. Ulmer Verlag, Stuttgart, 1051 S.
- OBERDORFER, E. (Hrsg.) (1993): Süddeutsche Pflanzengesellschaften. Wirtschaftswiesen und Unkrautgesellschaften. Teil III. 3. Aufl. Stuttgart, Fischer, 455 S.
- ORHAN, I., DELIORMAN-ORHAN, D. & B. ÖZÇELİK (2009): Antiviral activity and cytotoxicity of the lipophilic extracts of various edible plants and their fatty acids. *Food Chem.* 115: 701–705.
- PAUN, G., NEAGU, E., ALBU, C., MOROEANU, V. & G.-L. RADU (2016): Antioxidant activity and inhibitory effect of polyphenolic-rich extract from *Betonica officinalis* and *Impatiens noli-tangere* herbs on key enzyme linked to type 2 diabetes. *J. Taiwan Inst. Chem. Eng.* 60: 1–7.
- POTTS, S. G., BIESMEIJER, J. C., KREMEN, C., NEUMANN, P., SCHWEIGER, O. & W. E. KUNIN (2010): Global pollinator declines: Trends, impacts and drivers. *Trends Ecol. Evol.* 25 (6): 345–353.
- RAINE, N. E. & R. J. GILL (2015): Tasteless pesticides affect bees in the field. *Nature* 521: 3840.
- RAZALI, F. N., SINNIH, S. K., HUSSIN, H., ZAINAL ABIDIN, N. & A. S. SHUIB (2016): Tumor suppression effect of *Solanum nigrum* polysaccharide fraction on breast cancer via immunomodulation. *Int. J. Biol. Macromol.* 92: 185–193.
- SAUERWEIN, B. (2007): Säume in der Landnutzungsgeschichte. LEBBIMUK. Abhandl. Ber. Lebend. Bienenmuseum Knüllwald 4: 5–20.
- SCHEPER, J., REEMER, M., VAN KATS, R., OZINGA, W. A., VAN DER LINDEN, G. T. J., SCHAMINÉE, J. H. J., SIEPEL, H. & D. KLEIJN (2014): Museum specimens reveal loss of pollen host plants as key factors driving wild bee declines in The Netherlands. *PNAS* 111: 17555–17557.
- SCHUMACHER, W. (1980): Flora und Vegetation der Äcker, Raine und Ruderalplätze. Deutscher Naturschutzring, Bundesverband für Umweltschutz (Hrsg.). Bonn-Oberkassel.
- SENDL, A., MULINACCI, N., VINCIERI, F. F. & H. WAGNER (1993): Anti-inflammatory and immunologically active polysaccharide of *Sedum telephium*. *Phytochemistry* 34: 1357–1362.
- SENEJOUX, F., DEMOUGEOT, C., CUCIUREANU, M., MIRON, A., CUCIUREANU, R., BERTHELOT, A. & C. GIRARD-THERNIER (2013): Vasorelaxant effects and mechanisms of action of *Heracleum sphondylium* L. (*Apiaceae*) in rat thoracic aorta. *J. Ethnopharm.* 147: 536–539.
- SERKEDJIEVA, J. (1995): Inhibition of influenza virus protein synthesis by a plant preparation from *Geranium sanguineum* L. *Acta Virol.* 39: 5–10.
- SHARIFI, N., SOURI, E., ZIAI, S. A., AMIN, G. & M. AMANLOU (2013): Discovery of new angiotensin converting enzyme (ACE) inhibitors from medicinal plants to treat hypertension using an in vitro assay. *DARU J. Pharm. Sci.* 21(1): 1–8.
- SHEBABY, W. N., DAHER, C. F., EL-SIBAI, M., BODMAN-SMITH, K., MANSOUR, A., KARAM, M. C. & M. MROUEH (2015): Antioxidant and hepatoprotective activities of the oil fractions from wild carrot (*Daucus carota* ssp. *carota*). *Pharm. Biol.* 53: 1285–1294.
- SHU, X. S., LV, J. H., TAO, J., LI, G. M., LI, H. D. & N. MA (2009): Antihyperglycemic effects of total flavonoids from *Polygona-*

tum odoratum in STZ and alloxan-induced diabetic rats. J. Ethnopharmacol. 124: 539–543.

SIGURDSSON, S. & S. GUDBJARNASON (2007): Inhibition of acetylcholinesterase by extracts and constituents from *Angelica archangelica* and *Geranium sylvaticum*. Z. Naturforsch. C 62: 689–693.

SLAVÍK, J., ZÁCKOVÁ, P., MICHLOVÁ, J., OPLETAL, L. & M. SOVOVÁ (1994): Phytotherapeutic aspects of diseases of the circulatory system. III: Cardiotonic and cardiotoxic effects of hyrcanoside and deglucohyrcanoside isolated from *Coronilla varia* L. Ceska. Slov. Farm. 43: 298–302.

SOLOGUB, V. & A. GRYTSYK (2013): The research of the *Hypericum* extracts' pharmacological activity. Pharma Innovation 1: 85–89.

TABASSUM, N. & F. AHMAD (2011): Role of natural herbs in the treatment of hypertension. Pharmacogn. Rev. 5: 30–40.

TOVCHIGA, O.V. (2016): The influence of goutweed (*Aegopodium podagraria* L.) tincture and metformin on the carbohydrate and lipid metabolism in dexamethasone-treated rats. BMC Complement. Altern. Med. 16: 235–246.

VANBERGEN, A.J. & THE INSECT POLLINATORS INITIATIVE (2013): Threats to an ecosystem service: pressures on pollinators. Front. Ecol. Environ. 11: 251–259.

VLASE, L., BENEDEC, D., HANGANU, D., DAMIAN, G., CSILLAG, I., SEVASTRE, B., MOT, A.C., SILAGHI-DUMITRESCU, R. & I. TILEA (2014): Evaluation of antioxidant and antimicrobial activities and phenolic profile of *Hyssopus officinalis*, *Ocimum basilikum* and *Teucrium chamaedrys*. Molecules 19: 5490–5507.

YESILADA, E. & E. KÜPELİ (2007): *Clematis vitalba* L. aerial part exhibits potent anti-inflammatory, antinociceptive and antipyretic effects. J. Ethnopharmacol. 110: 504–515.

ZĂHAN, M., MICLEA, I., CRISTE, A. & V. MICLEA (2013): Modulation of murine macrophages phagocytotic activity by polysaccharide extract from *Chenopodium bonus henricus*. Centr. Eur. J. Immunol. 38: 183186.

Autorin

Apl. Prof. Dr. Mechthild Neitzke
 Arnulfstraße 35
 54295 Trier
 mechthild.neitzke@gmx.de